

Full unedited gel for Figure 4A, lanes used (1-5)



4-20% SDS-PAGE

Primary Ab: 1:1,000 rabbit anti-human C1-INH mAb (Abcam AB134918), secondary: 1:10,000 horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG (Abcam).


Full unedited gel for Figure 5A, lanes used (1-8)



4-20% SDS-PAGE

Primary Ab: 1:1,000 rabbit anti-human C1-INH mAb (Abcam AB134918), secondary: 1:10,000 horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG (Abcam).


Full unedited gel for Figure 6A, lanes used (1-8)



4-20% SDS-PAGE

Primary Ab: 1:1,000 rabbit anti-human C1-INH mAb (Abcam AB134918), secondary: 1:10,000 horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG (Abcam).


Full unedited gel for Figure 6B, lanes used (1-10)




Full unedited gel for Figure 7A, lanes used (1-9)

4-20% SDS-PAGE

Primary Ab: 1:1,000 rabbit anti-human C1-INH mAb (Abcam AB134918), secondary: 1:10,000 horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG (Abcam).



Full unedited gel for Figure 7B, lanes used (1-8)



4-20% SDS-PAGE

Primary Ab: 1:1,000 rabbit anti-human C1-INH mAb (Abcam AB134918), secondary: 1:10,000 horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG (Abcam).